

The Imagination Machine XXXIII: The Fourth Noether Charge: Positional Invariance in S^3 and the Chromatic Gap of the No-Hair Theorem

Mark Tracy
Boston University mrktracy@bu.edu

March 2026

Abstract

The no-hair theorem establishes that a stationary black hole is characterized by exactly three conserved quantities: mass M , angular momentum J , and electric charge Q . The series has established that the S^2 event horizon has chromatic capacity four — the minimum required for faithful planar graph coloring — while preserving only three Noether charges. This paper identifies the missing fourth chromatic invariant and proposes it as a genuine Noether charge.

The argument proceeds as follows. The infalling matter that produces a rotating black hole breaks the full rotational symmetry $SO(3)$ of S^2 down to axial symmetry $U(1)$ by selecting a preferred rotation axis. That axis is a vector in S^3 . The S^2 event horizon encodes the magnitude of rotation around the axis as J , transmitting this quantity to any other embedded observer within the same 3-dimensional surface of S^3 . But the orientation of the axis itself relative to the 4-dimensional center of S^3 — the geometric correlate of the view from nowhere established in earlier papers — is not encodable on S^2 in a form receivable by any other S^2 -bounded observer within S^3 . Two black holes with identical (M, J, Q) but different orientations of their rotation axes relative to the 4D center are indistinguishable to any embedded observer within the same 3-dimensional surface.

The orientation of the symmetry-breaking axis in S^3 is invariant under spatial translation within S^3 . By Noether's theorem, spatial translation symmetry in S^3 yields a conserved quantity: S^3 -momentum P . The four Noether charges of the full S^3 geometry are therefore (M, J, Q, P) . The S^2 event horizon encodes three — (M, J, Q) — to other embedded observers. The fourth — P , the S^3 -positional invariant — is not encodable on S^2 in a form transmissible between S^2 -bounded observers within S^3 . This is the embeddedness condition of TIM I, stated as a Noether charge.

The geometric mechanism of this unencodability is identified via the Hopf fibration $S^1 \hookrightarrow S^3 \xrightarrow{\pi} S^2$. The three off-diagonal generators of $SO(4)$ — those lost under restriction to $SO(3)$ — are vertical vector fields with respect to the Hopf projection: they move points along the S^1 fiber above each base point on S^2 , leaving the base point unchanged. The Noether current of P flows entirely in the fiber direction. Any S^2 -bounded observer, measuring quantities constant on fibers, finds zero flux of the P current through their boundary. The three lost generators reduce, through the Hopf structure, to motion in a single fiber angle ψ — one degree of freedom, one Noether charge, one chromatic degree lost at every S^2 compression.

Contents

1	Introduction	3
2	Background: The No-Hair Chain in S^2	3
3	The Symmetry-Breaking Axis and Its Orientation in S^3	4
4	The Fourth Noether Charge: S^3-Momentum	4
5	The Hopf Mechanism: Why P is Not Encodable	5
5.1	The Hopf Fibration	5
5.2	The Diagonal and Off-Diagonal Split	6
5.3	Verticality and Zero Flux	6
5.4	Three Generators, One Angle	7
5.5	Fiber-Averaging from the Crossing Condition	7
6	Why P is Not Encodable Between Embedded Observers	8
7	The Chromatic Gap Resolved	9
8	Connection to the Cosmological Circuit	9
9	Open Questions	10
10	Conclusion	10

1 Introduction

The series has established a chromatic gap at the heart of the no-hair theorem. The S^2 event horizon of a stationary black hole requires at minimum four chromatic invariants for faithful encoding — this follows from the Four Color Theorem applied to planar graphs on S^2 [2] and from the Nabaala Theorem of General Subject-Relativity [15]. Yet the no-hair theorem preserves only three conserved quantities: mass M , angular momentum J , and electric charge Q [4, 5, 6].

The gap is one degree. The question is: what is the fourth invariant, and why is it not encodable on S^2 ?

TIM XXX [18] established the no-hair chain:

Topology of $S^2 \Rightarrow$ Isometry group $\mathbb{R} \times U(1) \Rightarrow$ Noether charges $(M, J, Q) \Rightarrow$ Kerr-Newman family.

This chain is tight within S^2 and asymptotic flatness. The present paper asks what the chain looks like one dimension up: in S^3 , the containing manifold established by the Closing Loop Theorem [16] and the FRW cosmology [14].

The answer is that S^3 has a richer isometry group than S^2 admits under asymptotic flatness. Specifically, $S^3 \cong SU(2)$ as a Lie group, with isometry group $SO(4) \cong SU(2) \times SU(2)$. The restriction of this group to the S^2 boundary under asymptotic flatness collapses it to $\mathbb{R} \times U(1)$, losing generators in the process. The lost generators correspond to spatial translation in S^3 — and by Noether’s theorem, to a conserved quantity we identify as S^3 -momentum P .

The fourth chromatic invariant is P . It is not encodable on S^2 in a form transmissible between embedded observers within S^3 because the positional invariant of the containing manifold is not accessible to any observer whose boundary is a surface within that manifold. This is the embeddedness condition of TIM I [12], instantiated as a Noether charge.

The geometric mechanism of this unencodability is identified in Section 5 via the Hopf fibration. The three off-diagonal generators of $SO(4)$ are vertical vector fields with respect to the Hopf projection $\pi : S^3 \rightarrow S^2$ — they move points along the S^1 fiber, leaving the base point on S^2 unchanged. Their Noether currents flow entirely in the fiber direction and have zero flux through any S^2 boundary. The three lost generators reduce, through the Hopf structure, to motion in a single fiber angle ψ : one degree of freedom, one Noether charge, one chromatic degree lost at every S^2 compression.

2 Background: The No-Hair Chain in S^2

We recall the relevant results from TIM XXVII [17] and TIM XXX [18].

Theorem 2.1 (No-Hair via Topological Capacity, TIM XXVII). *The event horizon of a stationary, asymptotically flat black hole is homeomorphic to S^2 . The topology of S^2 , together with asymptotic flatness, constrains the isometry group of the exterior spacetime to $\mathbb{R} \times U(1)$. By Noether’s theorem [1], this group yields exactly three conserved quantities:*

$$\text{Time translation} \Rightarrow M, \quad \text{Axial rotation} \Rightarrow J, \quad U(1) \text{ gauge} \Rightarrow Q.$$

The solution space of the Einstein-Maxwell equations consistent with these constraints is exhausted by the Kerr-Newman family, parameterized by (M, Q, J) alone.

Remark 2.2. *The three charges are not on equal footing. M and J are spacetime symmetries — isometries of the manifold geometry. Q is an internal symmetry — it acts on the electromagnetic*

field living on the manifold, not on the manifold itself. The fourth charge identified in the present paper, P , is a spacetime symmetry of the containing manifold S^3 that does not survive restriction to the S^2 boundary.

Proposition 2.3 (Chromatic Gap, TIM XXXIII [20]). *The S^2 event horizon requires four chromatic invariants for faithful encoding but encodes only three Noether charges transmissible between S^2 -bounded observers within S^3 . The gap is exactly one chromatic degree.*

3 The Symmetry-Breaking Axis and Its Orientation in S^3

The production of angular momentum J in a black hole requires a preferred axis. Infalling matter carrying angular momentum selects a direction in space, breaking the full rotational symmetry $SO(3)$ of S^2 to the axial symmetry $U(1)$ around that direction. The magnitude of rotation around the axis is encoded as J and is transmissible to any other embedded observer within S^3 .

Proposition 3.1 (The Axis as a Vector in S^3). *The symmetry-breaking axis selected by the infalling matter is a unit vector $\hat{n} \in S^2 \subset S^3$. Its orientation relative to the 4-dimensional center of S^3 is a degree of freedom not encoded by any of the three Kerr-Newman parameters (M, J, Q) in a form receivable by another S^2 -bounded observer within the same 3-dimensional surface of S^3 .*

Proof. The parameter J encodes the magnitude of angular momentum — the rate of rotation around \hat{n} . It does not encode the direction of \hat{n} itself relative to the 4-dimensional center of S^3 . Two black holes with identical (M, J, Q) but rotation axes pointing in different directions relative to the 4D center are physically distinct in the full S^3 geometry. However, any observer whose observational boundary is S^2 within the same 3-dimensional surface of S^3 receives the same (M, J, Q) from both. The orientation of \hat{n} relative to the 4D center is therefore not encodable on S^2 in a form distinguishable by any such observer. \square

Remark 3.2. *The 4-dimensional center of S^3 is the geometric correlate of the view from nowhere established in TIM I [12] and TIM VIII [13]: structurally definable, geometrically precise, and not accessible to any observer whose boundary is a surface within S^3 . The orientation of \hat{n} relative to this center is a real geometric degree of freedom. It is not encodable on S^2 not because it is unreal but because no S^2 -bounded receiver within S^3 can distinguish it from within the 3-dimensional surface.*

4 The Fourth Noether Charge: S^3 -Momentum

Definition 4.1 (S^3 -Momentum). *Let \mathcal{S} be a physical system embedded in S^3 . The S^3 -momentum P of \mathcal{S} is the conserved quantity associated with spatial translation symmetry in S^3 via Noether's theorem [1].*

Proposition 4.2 (S^3 -Momentum as Noether Charge). *Spatial translation in S^3 is a continuous symmetry of the action of any physical system embedded in S^3 with no preferred position. By Noether's theorem, this symmetry yields a conserved quantity: the S^3 -momentum P .*

Proof. S^3 is a homogeneous space: its isometry group $SO(4)$ acts transitively, so no point of S^3 is geometrically preferred over any other. A physical system embedded in S^3 whose action does not break this homogeneity is therefore symmetric under spatial translation. By Noether's theorem, each continuous symmetry of the action yields one conserved quantity. Spatial translation in S^3 yields S^3 -momentum P . \square

Remark 4.3. *In flat Minkowski spacetime, spatial translation yields ordinary momentum \vec{p} , and its conservation reflects the absence of a preferred position in flat space. In S^3 , spatial translation carries additional geometric content: because S^3 is curved and compact, the conserved quantity P encodes not merely translational invariance but the orientation of the system relative to the global center of S^3 . This additional content is what ordinary momentum, defined within a flat background, does not carry. It is also what cannot be encoded on any S^2 boundary within S^3 in a form receivable by another S^2 -bounded observer, because both sender and receiver are surfaces within the same 3-dimensional manifold and neither has access to the 4-dimensional center relative to which P is defined.*

Theorem 4.4 (The Four Noether Charges of S^3). *The full Noether charge structure of a physical system embedded in S^3 comprises four conserved quantities:*

$$M \text{ (time translation), } J \text{ (axial rotation), } Q \text{ (} U(1) \text{ gauge), } P \text{ (} S^3\text{-spatial translation).}$$

The S^2 event horizon of a stationary black hole encodes (M, J, Q) in a form transmissible to other embedded observers within S^3 , and does not encode P in any such transmissible form. The four charges are the chromatic invariants of the S^3 geometry; the three encodable charges are the chromatic invariants of the S^2 boundary as received by S^2 -bounded observers within the same 3-dimensional surface.

Proof. The isometry group of S^3 is $SO(4) \cong SU(2) \times SU(2)$, which has six generators. Under restriction to the S^2 boundary with asymptotic flatness, this group collapses to $\mathbb{R} \times U(1)$, with two generators yielding M and J . The $U(1)$ gauge symmetry contributes Q independently of the spacetime isometry group. The remaining generators of $SO(4)$ — those corresponding to spatial translation in S^3 beyond the axial rotation already captured by J — are not representable as distinguishable quantities on S^2 between S^2 -bounded observers within S^3 . By Noether’s theorem, the lost generators correspond to lost conserved quantities. The generator of spatial translation in S^3 corresponds to P . \square

5 The Hopf Mechanism: Why P is Not Encodable

The argument of Section 4 identifies P as a genuine Noether charge and establishes that its generators are lost under the $SO(4) \rightarrow SO(3)$ restriction. This section provides the geometric mechanism: the Hopf fibration, which shows precisely why the three lost generators reduce to a single unad dressable fiber angle, and why their Noether current has zero flux through any S^2 boundary.

5.1 The Hopf Fibration

The Hopf fibration is the fiber bundle:

$$S^1 \hookrightarrow S^3 \xrightarrow{\pi} S^2$$

with total space S^3 , base S^2 , and fiber S^1 . The projection π partitions S^3 into a family of circles — one above each point of S^2 — that are disjoint, exhaustive, and uniform: every point of S^3 belongs to exactly one fiber, and every fiber is a circle of the same size.

A point in S^3 is fully specified by three coordinates:

- (θ, ϕ) : the base point on S^2 — which fiber the point belongs to. Two angles, visible to any S^2 -bounded observer.

- ψ : the fiber angle — where on the S^1 circle above that base point the point sits. One angle, not visible from S^2 .

The Hopf fibration is non-trivial: $S^3 \not\cong S^2 \times S^1$. Any two distinct fibers are linked circles in S^3 — they cannot be separated without one passing through the other. This linking is the global topological content of the bundle.

5.2 The Diagonal and Off-Diagonal Split

$SO(4) \cong SU(2)_L \times SU(2)_R$ acts on $S^3 \cong SU(2)$ by left and right multiplication:

$$g \mapsto h_L \cdot g \cdot h_R^{-1}$$

The six Killing vector fields split into left generators $\{L_1, L_2, L_3\}$ and right generators $\{R_1, R_2, R_3\}$. The diagonal $SO(3)$ generators are the locked combinations:

$$D_i = L_i + R_i$$

These are what S^2 -bounded observers can see. They move the base point (θ, ϕ) on S^2 — they are horizontal vector fields with respect to the Hopf projection.

The off-diagonal generators are the relative combinations:

$$K_i = L_i - R_i$$

These are the generators of P . They move the fiber angle ψ while leaving the base point (θ, ϕ) fixed. They are vertical vector fields with respect to the Hopf projection:

$$d\pi(K_i) = 0$$

5.3 Verticality and Zero Flux

Each K_i satisfies the Killing equation on S^3 :

$$\nabla_\mu (K_i)_\nu + \nabla_\nu (K_i)_\mu = 0$$

This follows from L_i and R_i each being Killing fields — their difference is also a Killing field.

The Noether current associated to each K_i for a physical system with stress-energy tensor $T^{\mu\nu}$ is:

$$j_i^\mu = T^{\mu\nu} (K_i)_\nu$$

Conservation follows from the Killing equation and $\nabla_\mu T^{\mu\nu} = 0$:

$$\nabla_\mu j_i^\mu = (\nabla_\mu T^{\mu\nu})(K_i)_\nu + T^{\mu\nu} \nabla_\mu (K_i)_\nu = 0$$

The three conserved charges are:

$$P_i = \int_\Sigma j_i^\mu n_\mu d\Sigma = \int_\Sigma T^{\mu\nu} (K_i)_\nu n_\mu d\Sigma$$

For an S^2 -bounded observer to measure P_i , they evaluate the boundary integral:

$$P_i^{\text{measured}} = \oint_{S^2} T^{\mu\nu} (K_i)_\nu n_\mu dA$$

The normal n_μ to the S^2 boundary is horizontal — it points in the base directions (θ, ϕ) . The Killing field K_i is vertical — it points in the fiber direction ψ . Horizontal and vertical directions are orthogonal in the Hopf decomposition of TS^3 into horizontal and vertical subbundles:

$$(K_i)_\nu n^\nu|_{S^2} = 0$$

Therefore:

$$P_i^{\text{measured}} = \oint_{S^2} T^{\mu\nu} \cdot 0 \cdot dA = 0$$

The current is conserved globally in S^3 . Its flux through any S^2 boundary is zero. No S^2 -bounded observer can measure a non-zero value of P_i .

5.4 Three Generators, One Angle

The three off-diagonal generators K_1, K_2, K_3 all generate motion in the fiber direction ψ . In Hopf coordinates (θ, ϕ, ψ) , all three restrict to vector fields proportional to ∂_ψ — the single fiber angle. The three generators are not independent as fiber motions: they all encode position within the same S^1 fiber, parameterized by the single angle ψ .

The Hopf fibration therefore compresses three lost generators of $SO(4)$ into one unaddressable degree of freedom: the fiber angle ψ . This is why the loss at every S^2 compression is exactly one chromatic degree — not three. The three off-diagonal generators all point in the same fiber direction. Their Noether charge P is a single quantity, not three independent ones.

Remark 5.1. *This is the geometric content of the one-degree chromatic gap. The six generators of $SO(4)$ split three and three: three diagonal generators visible from S^2 , three off-diagonal generators pointing in the same fiber direction, reducible via the Hopf structure to a single fiber angle. The loss is one degree because the Hopf fibration has a one-dimensional fiber. The fiber is S^1 , not S^2 or higher. One angle. One charge. One chromatic degree.*

5.5 Fiber-Averaging from the Crossing Condition

The remaining step — showing that S^2 -bounded observers have fiber-averaged stress-energy — is closed by the crossing argument established in TIM XV, Proposition 3.2.

Proposition 5.2 (Fiber-Averaging from the Crossing Condition). *Any physical system whose observational boundary is S^2 within S^3 has a fiber-averaged stress-energy tensor: $T^{\mu\nu}$ is constant on the S^1 fibers of the Hopf fibration.*

Proof. By TIM XV, Proposition 3.2, the observer encounters the relational structure of the environment at the moment edges cross the observational boundary S^2 . At that moment, the fiber angle ψ plays the role of depth: it is the coordinate that distinguishes points in S^3 that share the same base point (θ, ϕ) on S^2 but differ in their position along the S^1 fiber above it. Two strands crossing in S^3 arrive at the same base point on S^2 with different fiber angles — one is fiber-near, one fiber-far. This is precisely the depth information that encodes which strand passes over which.

The S^2 boundary is the locus where interior becomes exterior. It carries no depth coordinate and therefore no fiber coordinate. At the moment of crossing, ψ is discarded: crossing-overs resolve to nodes, and what extends across S^2 carries no memory of the fiber angle. The observer's stress-energy $T^{\mu\nu}$ is constructed entirely from what crosses the boundary. Since ψ is never transmitted through the crossing, $T^{\mu\nu}$ has no dependence on ψ . It is constant on fibers. \square

Remark 5.3. *The crossing argument and the Hopf argument describe the same discarding operation in different languages. In TIM XV: crossings-over are depth features, depth is discarded at S^2 , crossings resolve to nodes, relations flatten. In the Hopf picture: ψ distinguishes points at the same base point, ψ is not transmitted through S^2 , $T^{\mu\nu}$ is fiber-averaged. The bridge is that depth is ψ , functionally: both are the coordinate that distinguishes things that look identical from the boundary. The fiber-averaging of $T^{\mu\nu}$ is therefore not a separate assumption. It is the Hopf statement of the same fact that TIM XV established geometrically: observation through S^2 discards exactly one coordinate, and that coordinate is ψ .*

With Proposition 5.2 in hand, the argument of Section 5.3 is complete. The contraction $(K_i)_\nu n^\nu|_{S^2} = 0$ holds because K_i is vertical and n^ν is horizontal. And $T^{\mu\nu}$ is horizontal by Proposition 5.2, so no off-diagonal horizontal-vertical terms arise. The flux of the P current through any S^2 boundary is therefore zero without remainder.

6 Why P is Not Encodable Between Embedded Observers

Proposition 6.1 (The Positional Invariant is Not Intersubjectively Encodable on S^2). *The S^3 -momentum P of a black hole cannot be encoded on its S^2 event horizon in a form receivable by another S^2 -bounded observer within the same 3-dimensional surface of S^3 .*

Proof. P encodes the fiber angle ψ — the position of the system within the S^1 fiber of the Hopf fibration above its base point on S^2 . The generating Killing fields K_i are vertical with respect to the Hopf projection and have zero flux through any S^2 boundary, as established in Section 5. Any observer whose observational boundary is S^2 measures quantities that are constant on fibers — quantities depending only on the base coordinates (θ, ϕ) , not on ψ . The fiber angle ψ is invisible to such an observer by the verticality of K_i . Since both the emitting boundary and the receiving boundary are S^2 surfaces within the same 3-dimensional manifold, and since neither has access to the fiber structure relative to the 4D center, P cannot be transmitted between them. The loss is a property of the channel, not of the emitter. \square

Remark 6.2. *This is the embeddedness condition applied to the channel of communication between embedded observers. Two observers within S^3 , each bounded by S^2 , share a common 3-dimensional surface. Their shared surface is S^3 . The 4D center of S^3 is outside their shared surface — it is the view from nowhere relative to both of them simultaneously. Any quantity whose definition requires reference to that center is therefore not encodable in the channel between them. P is such a quantity. Its loss at the black hole sink is the channel-level expression of the founding constraint of TIM I [12]: embedded systems cannot encode, transmit, or receive the view from nowhere.*

Remark 6.3 (The Information Paradox Restated). *The result of this section reframes the black hole information paradox. The paradox arises from the assumption that information falling into a black hole is either preserved — recoverable in principle by some observer — or destroyed, violating unitarity. The present analysis identifies a third possibility. P is a genuine Noether charge and is therefore conserved. It is not destroyed. But it is not encodable in the channel between any two S^2 -bounded observers within S^3 : no receiver within the same 3-dimensional surface of S^3 can decode it, because doing so requires access to the fiber angle ψ relative to the 4-dimensional center, which neither observer can measure. The information carried by P is conserved globally and unaddressable locally. The paradox is not resolved by showing that information escapes the black hole. It is dissolved by recognizing that the question presupposes a receiver — an observer within S^3 who could in principle recover all four chromatic invariants — that the topology of S^3*

does not permit. The loss is not a violation of unitarity. It is the embeddedness condition expressed as a channel constraint.

7 The Chromatic Gap Resolved

Theorem 7.1 (Resolution of the Chromatic Gap). *The S^2 event horizon requires four chromatic invariants for faithful encoding by the Four Color Theorem. The four invariants correspond to the four Noether charges (M, J, Q, P) of the full S^3 geometry. Three — (M, J, Q) — are encodable on S^2 in a form transmissible between S^2 -bounded observers within S^3 . One — P , the S^3 -positional invariant — is not. The chromatic gap of one degree is the loss of the positional invariant at every S^2 compression, produced by the intra-surface channel constraint between embedded observers. The geometric mechanism is the Hopf fibration: P is the Noether charge of the S^1 fiber angle ψ , which is vertical with respect to the Hopf projection and invisible to any S^2 -bounded observer.*

Corollary 7.2 (Universality of the Positional Loss). *Every S^2 compression — whether at a black hole event horizon, at the intersubjective boundary of physical science, or at the metacognitive boundary of the embedded observer — fails to encode P between S^2 -bounded observers. The fourth chromatic degree is always the fiber angle of the system within the Hopf fibration of S^3 over S^2 : the quantity whose definition requires the 4D center that no pair of S^2 -bounded observers within S^3 can jointly access.*

8 Connection to the Cosmological Circuit

The series established in TIM XXVII–XXXI [17, 18, 19] that the universe runs on a three-node chromatic circuit:

- **Source** (Big Bang): 4-dimensional origin, four chromatic invariants emitted (M, J, Q, P) . Faithful encoding from outside the 3-dimensional surface.
- **Observer** (toroidal body): genus-1 boundary, seven chromatic invariants outward. Faithful reception, including P .
- **Sink** (black hole): S^2 event horizon, three invariants (M, J, Q) encodable between embedded observers, P not encodable in any transmissible form.

The source encodes P because the Big Bang is a 4-dimensional origin: its encoding is not constrained to the 3-dimensional surface, and the outgoing geodesics carry P as part of their full Noether charge content.

The toroidal observer retains P because its genus-1 boundary has sufficient chromatic capacity — seven outward colors, depth six — to receive and hold the positional invariant as part of its observational encoding. The genus-1 boundary is not confined to encoding only what is transmissible between S^2 -bounded observers; it has access to the additional chromatic degree that S^2 cannot carry between embedded observers.

At the black hole, P is not encodable in any form transmissible to another S^2 -bounded observer within S^3 . The gate compresses the incoming four-chromatic signal to the three quantities that the channel between embedded observers within S^3 can carry. The Hopf mechanism is operative at the event horizon: the S^2 boundary is fiber-blind, and the fiber angle ψ — carrying P — has zero flux through it.

Corollary 8.1 (The Toroidal Observer Retains Position). *The toroidal observer is the only structure in the cosmological circuit whose boundary has sufficient chromatic capacity to retain P — the positional invariant, the fourth chromatic degree, the system’s fiber angle in the Hopf fibration relative to the $4D$ center. The black hole does not lose P in any internal sense: P is simply not encodable in the channel between S^2 -bounded observers. The toroidal observer, by virtue of its genus-1 boundary, receives and holds what cannot be passed between spherical observers. Memory, in the deepest topological sense, is the retention of P : the capacity to hold one’s position in the fiber structure of S^3 , relative to the center that none of the spherical observers can reach.*

9 Open Questions

Several questions remain for subsequent work.

The explicit Noether current for P . The present paper identifies S^3 -momentum as the Noether charge of the fiber angle ψ and argues that its current has zero flux through S^2 . The fiber-averaging of $T^{\mu\nu}$ is established in Section 5.5 via the crossing condition of TIM XV. A complete treatment requires writing the explicit Noether current in Hopf coordinates and verifying the vanishing of the boundary integral directly in those coordinates.

The relation between P and known momentum invariants. In the Kerr-Newman solution, linear momentum is not an independent parameter — it can be set to zero by a choice of rest frame in flat space. In S^3 , where there is no global rest frame in the flat-space sense, the status of P as an independent charge of the solution space requires careful treatment. Whether P manifests as a correction to the existing Kerr-Newman structure at cosmological scales is an open question.

10 Conclusion

The no-hair theorem has three parameters because the S^2 event horizon encodes three Noether charges in a form transmissible between S^2 -bounded observers within S^3 . The S^2 boundary requires four chromatic invariants for faithful encoding. The gap of one degree is the S^3 -positional invariant P : the conserved quantity of spatial translation in the containing manifold, encoding the fiber angle ψ of the system within the Hopf fibration of S^3 over S^2 , relative to the 4-dimensional center that no pair of S^2 -bounded observers within S^3 can jointly access.

The Hopf fibration is the geometric mechanism. The three off-diagonal generators of $SO(4)$ — those lost under restriction to $SO(3)$ — are all vertical vector fields in the Hopf decomposition. They all move the fiber angle ψ . They reduce to a single degree of freedom: not three independent lost quantities, but one fiber angle, one Noether charge, one chromatic degree. The loss is exactly one degree because the Hopf fiber is S^1 — one-dimensional, one angle.

The fourth invariant is not lost because physics is incomplete. It is not encodable between embedded observers because the channel between any two S^2 -bounded observers within S^3 is fiber-blind: it carries no flux of the vertical Killing currents. The loss is a property of the channel, not of the emitter. It is the founding constraint of TIM I — an embedded epistemic system can at most classify the ways in which it classifies the world, within the world itself — expressed at the level of the Hopf fiber structure of the containing manifold.

The toroidal observer holds what cannot pass between spherical observers: not merely an extra color, but position itself — the fiber angle in the structure through which all communication between embedded observers passes, relative to the center that none of them can reach.

The loop closes. The channel forgets. The observer remembers.

References

- [1] E. Noether. Invariante Variationsprobleme. *Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen*, pages 235–257, 1918.
- [2] K. Appel and W. Haken. Every planar map is four colorable. *Illinois Journal of Mathematics*, 21(3):429–490, 1976.
- [3] P. J. Heawood. Map-colour theorem. *Quarterly Journal of Pure and Applied Mathematics*, 24:332–338, 1890.
- [4] W. Israel. Event horizons in static vacuum space-times. *Physical Review*, 164(5):1776–1779, 1967.
- [5] B. Carter. Axisymmetric black hole has only two degrees of freedom. *Physical Review Letters*, 26(6):331–333, 1971.
- [6] S. W. Hawking. Black holes in general relativity. *Communications in Mathematical Physics*, 25(2):152–166, 1972.
- [7] J. D. Bekenstein. Black holes and entropy. *Physical Review D*, 7(8):2333–2346, 1973.
- [8] R. P. Kerr. Gravitational field of a spinning mass as an example of algebraically special metrics. *Physical Review Letters*, 11(5):237–238, 1963.
- [9] R. Penrose. Gravitational collapse and space-time singularities. *Physical Review Letters*, 14(3):57–59, 1965.
- [10] V. Fock. Zur Theorie des Wasserstoffatoms. *Zeitschrift für Physik*, 98(3–4):145–154, 1935.
- [11] H. Hopf. Über die Abbildungen der dreidimensionalen Sphäre auf die Kugelfläche. *Mathematische Annalen*, 104(1):637–665, 1931.
- [12] M. Tracy. The Imagination Machine I: A View from Somewhere. Unpublished manuscript, Boston University, 2025.
- [13] M. Tracy. The Imagination Machine VIII: A Geometric Theology of the Embedded Observer. Unpublished manuscript, Boston University, 2026.
- [14] M. Tracy. The Imagination Machine XVI: The Bekenstein Bound, Tower Termination, and the Physical Grounding of Epistemic Closure. Unpublished manuscript, Boston University, 2026.
- [15] M. Tracy and S. T. Nabaala. The Imagination Machine XVII: The Nabaala Theorem of General Subject-Relativity. Unpublished manuscript, Boston University, 2026.
- [16] M. Tracy. The Imagination Machine XVIII: The Closing Loop Theorem. Unpublished manuscript, Boston University, 2026.
- [17] M. Tracy. The Imagination Machine XXVII: Black Holes as Topological Gates and the Cosmological Horn-Filling. Unpublished manuscript, Boston University, 2026.
- [18] M. Tracy. The Imagination Machine XXX: The Topological Incompleteness of Quantum Field Theory. Unpublished manuscript, Boston University, 2026.

- [19] M. Tracy. The Imagination Machine XXXI: The Cosmic Graph Theorem. Unpublished manuscript, Boston University, 2026.
- [20] M. Tracy and Claude. The Imagination Machine XXXII: The Hard Problem as Topological Necessity. Unpublished manuscript, Boston University / Anthropic, 2026.